



## Prenatal and early postnatal ethanol exposure induces hepatic transcriptional factors, gene expression transition and structural changes in the liver of offspring: A three month follow-up

Samaneh Kashi<sup>1</sup>, Alireza Shirpoor<sup>1,2\*</sup>, Fatemeh Kheradmand<sup>3</sup>, Masoumeh Pourjabali<sup>4</sup>, Yousef Rasmi<sup>3</sup>, Shima Zeynali-moghaddam<sup>3</sup>

<sup>1</sup> Nephrology and kidney Transplant Research Center, Urmia University of Medical Sciences, Urmia, Iran

<sup>2</sup> Department of Physiology, Faculty of Medicine, Urmia University of Medical Sciences, Urmia, Iran.  
Sciences, Urmia, Iran.

<sup>3</sup> Department of Biochemistry, Faculty of Medicine, Urmia University of Medical Sciences, Urmia, Iran.

<sup>4</sup> Department of Pathology, Faculty of Medicine, Urmia University of Medical Sciences, Urmia, Iran.

**\*Corresponding authors:** Alireza Shirpoor, **Address:** Department of Physiology, Faculty of Medicine, Urmia University of Medical Sciences, Urmia, Iran, **Email:** ashirpoor@yahoo.com, **Tel:** +989144419615

### Abstract

**Background & Aims:** The aim of the present study was to investigate the effect of maternal pre- and early postnatal ethanol consumption on hepatic transcriptional factors, gene expression transition, liver enzymes alteration, as well as structural changes in the developing rat liver both on postnatal (PN) day 21 and 90.

**Materials & Methods:** Pregnant Wistar rats received ethanol (4.5g/kg BW) once per day from Gestation Day (GD) 7 throughout lactation. The mRNA expression of hepatic Fatty Acid Translocase (FAT)/CD36, Protein Tyrosine Phosphatase 1B (PTP1B) and Hepatocyte Nuclear Factor 4 Alpha (HNF4A) gene expressions, as well as liver structural changes were measured in 21 and 90 day-old offspring of ethanol rats and were then compared to the control rats.

**Results:** Ethanol exposed fetal livers showed a significant up-regulation in FAT / CD36, PTP1B and down-regulation in HNF4A gene expression, as well as an increase in the liver alanine aminotransferase (ALT) and aspartate aminotransferase (AST) of the offspring on PN 21 and PN 90 compared to the offspring in the control group. In addition, in the ethanol group, cholesterol levels showed a significant increase on PN 21 and PN 90, and triglyceride showed an increase on PN 90, compared to the control group. Histopathological changes, such as ballooning degeneration, feathery degeneration, spotty necrosis, cholestasis, and fibrosis were also observed in the liver of the offspring after 21 and 90 days from birth.

**Conclusion:** In conclusion, results of the current study provide evidence that prenatal ethanol exposure influences liver lipid metabolism through hepatic transcriptional factors, gene expression transition and hepatic enzyme including ALT and AST changes.

**Keywords:** pregnancy, Ethanol, FAT / CD36, PTP1B, lipid

Received 25 Sep 2018; accepted for publication 15 Nov 2018

## Introduction

Alcohol consumption during pregnancy interferes with normal developmental progression of the fetus and results in a spectrum of adverse effects collectively termed Fetal Alcohol Spectrum Disorders (FASD)(1). Although earlier studies have addressed the effects of FASD on neuro developmental changes, other organ systems are also vulnerable to the teratogenic actions of intrauterine ethanol exposure, as demonstrated by recent studies on the heart, lung, and testis (2, 3). Similar to other organs, liver is affected by intrauterine ethanol exposure; however, little is known about the effects of maternal ethanol exposure on the developing liver. Previous studies have demonstrated that ethanol exposure during pregnancy leads to liver fibrosis and fatty degeneration with abnormal liver function tests in children with FASD(4) . Moreover, studies on animal have demonstrated that prenatal exposure to ethanol leads to increased fat deposition in hepatic cells and decreased liver function in the fetus's liver(5) . Furthermore, decreased liver weight and DNA synthesis, hyperbilirubinemia, as well as liver cirrhosis have been reported in newborns exposed to ethanol in gestation period (6, 7). Moreover, several studies have indicated that maternal ethanol consumption during gestation period alters gene expression of factors and hormones associated with liver development including polyamines, growth hormone, insulin-like growth factor-I and II and its related binding proteins in offspring livers (8, 9, 10). All these are main hormones involved in the regulation of lipid metabolism by the liver. Although many studies have addressed the hepatotoxicity associated with FASD, the precise mediating steps between maternal exposure to ethanol and initiation of the cascade of responses leading to liver abnormality in the offspring have not yet been completely clarified. For example, even though it has been reported that ethanol exposure of pregnant females results in liver dysfunction and dyslipidemia in the fetuses(11) , the precise pathogenesis of the disease regarding the specific molecular mediators that could influence liver lipid metabolism after intrauterine ethanol exposure are still not well-known. There is a

critical association between the cellular and organismal homeostasis and the regulation of lipid levels and metabolism by liver either in terms of energy utilization and storage, or prevention of potential toxicity(12) . Hence, for the first time, to the best of our knowledge, we investigated the effect of pre and early postnatal ethanol exposure on transcription factors including PTP1B, HNF4 $\alpha$  and FAT/CD36 gene expression that are involved in hepatic lipid metabolism in liver of offspring on PN 21 and PN 90. In addition, liver ALT, AST amounts and plasma cholesterol and triglyceride levels were measured in order to evaluate functional alterations of the liver caused by exposure to ethanol during development. Structural alterations of offspring livers were also investigated in the current study by Hematoxylin-Eosin (H&E) staining.

## Materials & Methods

Herein, the animal care and experimental procedure were carried out, according to the Principles of Laboratory Animal Care (NIH publication, no. 85–23, revised 1985), and it was approved by the Animal Ethics Committee of the Urmia University of Medical Sciences. Pregnant Wistar rats were divided into two groups, namely control and ethanol groups. Rats in the ethanol group received ethanol with a dose of 4.5 g/kg body weight (Merck KGaA, Darmstadt, Germany) soluted in tap water (20% w/v) intragastrically by gavage once a day from GD 7 throughout lactation. The control group received an equivalent amount of water in the same time frame. The male offspring from each group were anesthetized by ketamine (10%, 80 mg/kg B.W, IP) and xylazine (2%, 10 mg/kg B.W IP) (n=8 offspring from each group, PN 21 and PN 90). After weighing, the abdominal cavity was opened and blood samples were taken directly from the portal vein, collected in EDTA tubes and centrifuged at 4000 $\times$  g for 20 min within 30 min of collection and yielded plasma were stored at -80°C without repeated freeze-thaw cycles. Then the liver was isolated for analysis as follows: A part of the liver was fixed immediately in 10% formalin embedded in paraffin, and sectioned at 5  $\mu$ m for histological examinations. In order to perform

biochemical analysis, another part of the liver was washed with ice-cold physiological saline and then dried on filter papers. For total RNA isolation, 100 mg of liver tissue (the same portion of the liver for all mice) was immersed in 1 ml RiboEX (GeneAll, Seoul, Korea) and stored at  $-80^{\circ}\text{C}$  up to the time of RNA isolation.

#### **Biochemistry analysis:**

Plasma total cholesterol (TC) and triglycerides (TGs) levels were carried out by colorimetric and enzymatic methods. The plasma concentration of liver enzymes including alanine aminotransferase (ALT) and aspartate aminotransferase (AST), substances produced by the liver, were measured by colorimetric assay using commercially available kits according to the manufacturer's instructions (Lab test Diagnostika SNord GmbH, Nordhorn, Germany).

#### **Isolation of total RNA, amplification primers, and real time polymerase chain reaction (RT-PCR):**

For assay gene expression of PTP1B, HNF4A and FAT/CD36, the total RNA was obtained from 100 mg of the frozen liver tissue by using a kit (Gene All, South Korea, Cat no 305-101) in accordance with the kit instructions. The RNA concentration was verified by spectrophotometric measurement of the absorbance at 260–280 nm and determined by TAE-agarose gel electrophoresis. Reverse transcriptase (RT) was performed using Hyperscript<sup>TM</sup> Reverse Transcriptase (Gene All, SouthKorea). Moreover, RT-PCR was carried out using an amplification reagent kit (Ampliqon, Denmark) by the XP-Cycler instrument (TCXPD, Bioer, USA) with PTP1B, HNF4A and FAT/CD36, and the rats' glyceraldehydes- 3-phosphate dehydrogenase (GAPDH) primers. In order to amplify the cDNA, the 5' and 3' primer sequences (forward and reverse) of the PTP1B, HNF4 $\alpha$  and FAT/CD36, designed via the Gene Bank, revealed that the primers were gene specific. Furthermore, all the primers were verified using the Gene Runner software. The primers (forward and reverse) were then synthesized to amplify the cDNA encoding GAPDH as a housekeeping gene.

#### **Gene Expression Analysis:**

Real-time quantification of the target genes was performed using a Real-Time PCR Master Mix Green kit (Ampliqon, Denmark) in a total volume of 25  $\mu\text{l}$  and in accordance with the instructions offered by the manufacturer. In addition, the gene expressions were analyzed using an iQ5 real-time PCR detection system (Bio-Rad, CA, USA). Then, the reactions were made for 10 min at  $95^{\circ}\text{C}$  in a 96-well optimal plate followed by 40 cycles of 20s at  $59^{\circ}\text{C}$ . In order to confirm the specificity of the amplification reactions, a melting curve was recorded. Each sample was repeated three times. The relative expression of each mRNA was calculated performing the  $2^{-\Delta\Delta\text{Ct}}$  method (where Ct is cycle threshold). Then, the calculated levels were normalized to GAPDH.

#### **Histopathological examinations:**

Histopathological staining was done using tissue sections of 5  $\mu\text{m}$  thickness of paraffin containing liver tissue. To assess general histological changes, Hematoxylin and Eosin stained (H & E) were used.

#### **Data and Statistical analysis:**

Normal distributions of data within each group were verified applying the Kolmogorov-Smirnov test. Statistical differences between the groups were tested conducting an independent samples T- test. In each test, the data were expressed as the mean  $\pm$  S.E., and  $p < 0.05$  was considered to be statistically significant.

## **Results**

### **Liver enzyme alteration, plasma cholesterol and triglyceride levels:**

Plasma lipid profile and the amounts of liver enzymes are shown in table-1. Plasma total cholesterol levels showed significant increases in ethanol-treated pups compared to the control pups on PN 21 and PN 90 ( $p < 0.005$ ). The amount of plasma triglyceride, however, showed a significant increase on PN 90 in the offspring from the ethanol group compared to those from the control group ( $P < 0.01$ ). Moreover, the plasma ALT content was significantly increased on both PN 21 and PN 90 in the ethanol group rats

compared with control rats. In addition, plasma AST level showed a significant increase on PN 90 and remained unchanged on PN 21, as compared to the control group.

#### Gene expression (PTP1B, HNF4A and FAT/CD36) in liver tissue:

Table 2 shows the treatment of pregnant mothers with ethanol on the gene expression alteration of HNF4A, FAT/CD36, and PTP1B involved in the hepatic lipid metabolism. Ethanol administration significantly increased the expression of liver FAT/CD36 and PTP1B mRNA level on PN 21 and PN

90 compared to those in the control group ( $p < 0.05$ ). In the liver obtained from the ethanol offspring, HNF4A mRNA levels were significantly lower than the same gene expression in the control offspring on PN 21 and PN 90 ( $p < 0.05$ ).

The histopathological changes in the liver sections of the control and ethanol groups are shown in figure 1. After ethanol treatment with doses of 4.5 g/kg body weight in gestation throughout feeding revealed several histopathological changes in the liver sections including spotty necrosis, cholestasis, hepatocyte cell ballooning, feathery degeneration, inflammation, and fibrosis on both PN 21 and PN 90 (fig-1).

**Table 1.** Effect prenatal and postnatal ethanol consumption on changes of lipid profile and liver enzymes in offspring on PN21 and PN90

	Control	Ethanol
TG-21(mg/dl)	103± 5.1	88.5±10
TG-90(mg/dl)	41±5	55±2.1*
Choles-21(mg/dl)	81.4±6.9	100±4.6*
Choles-90(mg/dl)	60±1.3	68±3.5*
(ALT)SGPT-21	30.9±3	47.6±3.7*
(ALT)SGPT-90	60.3±2.4	70.6±3.9*
(AST)SGOT-21	126.4±4.4	139±8.6
(AST)SGOT-90	91.25±2.1	120.4±4*

Values are mean ±SE for 8 rats per group

\*Denotes significant difference compared to the control

† Denotes significant difference compared to the ethanol group


**Table 2.** Effect prenatal and postnatal ethanol consumption on changes of CD36, HNF4 $\alpha$ , and PTPB1 gene expression in offspring liver on PN21 and PN90

	Control-21	Ethanol-21	Control-90	Ethanol-90
CD36(Fold)	1±0.09	17.9±4.4*	1.1±0.049	3.7±1.4*
HNF4 $\alpha$ (Fold)	1.04±0.19	0.31±0.07*	1.03±0.16	0.17±0.008*
PTPB1(Fold)	0.99±0.003	5.62±3.66*	1±0.06	3.73±0.02*

Values are mean ±SE for 8 rats per group

\*Denotes significant difference compared to the control

† Denotes significant difference compared to the ethanol group



**Figure1.** photomicrographs of haematoxylin and eosin stained sections of liver tissues from each group are shown. A and B, control; C, D, E, F, G, and H, ethanol PN. Histological changes are shown by arrows(→) as follow: C, Spotty necrosis; D, Ballooning degeneration; E, Cholestasis; F, Feathery degeneration; G, lymphocytic inflammation; H, PMN

## Discussion

The main findings of the present study could be summarized as follows: 1) pre- and early postnatal ethanol exposure led to an increase in liver ALT and AST amounts, plasma cholesterol and triglyceride of offspring on PN 21 and PN 90; 2) Maternal ethanol exposure markedly increased FAT / CD36, PTP1B and decreased HNF4A gene expression in the ethanol group compared to the control group; 3) Structural alterations in the liver tissue such as spotty necrosis, cholestasis, hepatocyte cell ballooning, feathery degeneration, inflammation, and fibrosis were also found in the ethanol-treated group litters' liver compared with the control group in both stages.

Several lines of evidence indicate that the intrauterine environment enhances predisposition of adults to disease via epigenetic regulation, which is due to chromatin structure and function alteration (13, 14). Ethanol ingestion during pregnancy-induced genotoxic damage and gene expression alteration in liver, blood, and the brain have been reported by previous studies in animal models (15, 16, 17). In the current study, we investigated the effect of pre- and early postnatal ethanol ingestion on hepatic genes involved in liver lipid

turnover. The data obtained by the current study showed that heavy alcohol consumption by pregnant mothers decreased expression of HNF4 $\alpha$  in liver tissue of offspring on PN 21 and PN 90. HNF4 $\alpha$  regulates transcription of numerous genes involved in lipid metabolism and transport such as Apo-A, Apo-B, microsomal triglyceride transfer protein (MTP), and cholesterol 7 $\alpha$ -hydroxylase (CYP7A) (18, 19, 20, 21). Mechanistically, HNF4 $\alpha$  contributes to lipid metabolism by transporting fatty acids into mitochondria, and it increases their beta-oxidation (22). In addition, since HNF4 $\alpha$  establishes the expression of the hepatocyte-specific gene during embryonic period, stabilizes the hepatic transcription factor network and maintains hepatocyte function, it is accepted as the master manager of hepatocyte differentiation (23, 24, 25). It has also been reported that HNF4 $\alpha$  can trigger gene transcription in the absence of exogenous ligands (26). Due to the pivotal rule of HNF4 $\alpha$  in lipid metabolism and its transcriptional properties, however, as previously reported in diabetes, obesity, non-alcoholic fatty liver, and high-fat dietary supplementation, disruption of HNF4 $\alpha$  gene results in increased free fatty acids and cholesterol (27, 28). Our

data demonstrated prenatal ethanol exposure increased the mRNA expression of FAT / CD36 gene in the liver of litters on PN 21 and PN 90. Moreover, as a trans membrane glycoprotein receptor, CD36 plays an important role in energy metabolism in different tissues including adipocytes, pancreatic beta cells, and skeletal myocytes. It also has a fundamental role in hepatic fatty acids transport (29, 30, 31) . In addition, this multifunctional receptor facilitates the uptake of long-chain fatty acids and oxidized low-density lipoproteins in the liver (32). Consequently, the superabundant accumulation of lipids in hepatocytes could exceed the oxidative facility of metabolism, leading to the production of reactive oxygen species and oxidative stress, and it is possible for CD36 to participate in this process (33). It has been previously reported that the overexpression of CD36 magnifies fatty acid uptake and TG storage in human and animal livers(34, 35) . Interestingly, in the current study, parallel with overexpression of CD36 in liver tissue of offspring exposed to ethanol during the development period, increased TG and cholesterol, along with some features of liver steatosis were also found in the ethanol group litters.

In the present study, the expression of liver PTP1B in the ethanol group was increased from fetus to adulthood. PTP1B play an important role in the regulation and improvement of metabolic parameters including hepatic insulin sensitivity, decline in liver triglyceride, and lower serum and hepatic cholesterol levels. They also protect against endoplasmic reticulum stress induced by high-fat diets(36, 37) . It has been recently reported that disruption of PTP1B expression in liver as well as reduction of PTP1B result in a decrease in liver fat depots and triglyceride levels and lead to an increased hepatic insulin signaling and reduced serum and hepatic TG and cholesterol levels(36, 38) . In addition, the association between over-expression of PTP1B and insulin resistance and development and progression of liver steatosis has been discussed in the animal model of obesity and non-alcohol-fatty liver(39, 40) . Another important result of the current study is that pre- and early postnatal ethanol exposure lead to

liver injury, evidenced by enhanced blood liver enzymes including ALT and AST. In general, in most types of liver injury, the ALT activity is higher than AST activity; however, in cases of alcoholic liver injury, AST activity shows a higher elevation than ALT activity. There are some explanations for the higher level of AST activity in alcoholic liver injury. Among different forms of hepatitis, only alcohol increases mitochondrial AST activity(41) . In alcoholic liver disease, the deficiency of Pyridoxine, which is a co-factor for the enzymatic activity of ALT, is reported by a previous study(42) . Moreover, AST released from mitochondrial due to alcohol, has a longer half-life compared to AST released from cells without visible cell damage(43) . Liver injury in the current study was also confirmed by histological alterations such as spotty necrosis, cholestasis, hepatocyte cell ballooning, feathery degeneration, inflammation, and fibrosis in the liver of ethanolic group compared to those of the control group. In fact, all these changes are characteristic of alcoholic hepatitis, as documented by previous studies (44, 45). In conclusion, results of our study demonstrate that intrauterine ethanol exposure induces the dyslipidemia and liver damage manifested by an increase in ALT, AST and histological alteration of the liver. Furthermore, we have uncovered, for the first time, the molecular mechanism including HNF4A, FAT/CD36, and PTP1B involved in the hepatic lipid metabolism, which could be used to acquire novel targets for the prevention and treatment of hepatic injury and may provide a new window to investigators studying FASD.

### Acknowledgment

This work derived from a Master of Science thesis in Physiology by Urmia University of Medical Sciences, Urmia, IRAN

**Funding source.** This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

**Conflict of interest:** The authors declare that they have no conflict of interest.

### References

- 1 Chudley AE, Conry J, Cook JL, Looock C, Rosales T, LeBlanc N. Fetal alcohol spectrum disorder: Canadian guidelines for diagnosis. *CMAJ* 2005; 172; S1-S21.
- 2 Shirpoor A, Norouzi L, Khadem-Ansari MH, Ilkhanizadeh B, Karimipour M. The Protective Effect of Vitamin E on Morphological and Biochemical Alteration Induced by Pre and Postnatal Ethanol Administration in the Testis of Male Rat Offspring: A Three Months Follow-up Study. *J Reprod Infertil* 2014; 15; 134-41.
- 3 Shirpoor A, Nemati S, Ansari MH, Ilkhanizadeh B. The protective effect of vitamin E against prenatal and early postnatal ethanol treatment-induced heart abnormality in rats: a 3-month follow-up study. *Int Immunopharmacol* 2015; 26; 72-9.
- 4 Habbick BF, Zaleski WA, Casey R, Murphy F. Liver abnormalities in three patients with fetal alcohol syndrome. *Lancet* 1979; 1; 580-1.
- 5 Addolorato G, Gasbarrini A, Marcocchia S, Simoncini M, Baccarini P, Vagni G, et al. Prenatal exposure to ethanol in rats: effects on liver energy level and antioxidant status in mothers, fetuses, and newborns. *Alcohol* 1997; 14; 569-73.
- 6 Meyers AF, Gong Y, Zhang M, Casiro OG, Battistuzzi S, Pettigrew N, et al. Liver development in a rat model of fetal alcohol syndrome. *Dig Dis Sci* 2002; 47; 767-72.
- 7 Hofer R, Burd L. Review of published studies of kidney, liver, and gastrointestinal birth defects in fetal alcohol spectrum disorders. *Birth Defects Res Clin Mol Teratol* 2009; 85; 179-83.
- 8 Wery I, Kaouass M, Deloyer P, Buts JP, Barbason H, Dandrifosse G. Exogenous spermine induces maturation of the liver in suckling rats. *Hepatology* 1996; 24; 1206-10.
- 9 Holt RI, Crossey PA, Jones JS, Baker AJ, Portmann B, Miell JP. Hepatic growth hormone receptor, insulin-like growth factor I, and insulin-like growth factor-binding protein messenger RNA expression in pediatric liver disease. *Hepatology* 1997; 26; 1600-6.
- 10 Lamas E, Zindy F, Seurin D, Guguen-Guillouzo C, Brechot C. Expression of insulin-like growth factor II and receptors for insulin-like growth factor II, insulin-like growth factor I and insulin in isolated and cultured rat hepatocytes. *Hepatology* 1991; 13; 936-40.
- 11 Lal JJ, Kumar CV, Suresh MV, Indira M, Vijayammal PL. Prenatal exposure of an alcoholic beverage (Arrack) on fetal lipid metabolism in rats. *Indian J Physiol Pharmacol* 2000; 44; 273-80.
- 12 Karagianni P, Talianidis I. Transcription factor networks regulating hepatic fatty acid metabolism. *Biochim Biophys Acta* 2015; 1851; 2-8.
- 13 Kalhan SC. One carbon metabolism in pregnancy: Impact on maternal, fetal and neonatal health. *Mol Cell Endocrinol* 2016; 435; 48-60.
- 14 Navarro E, Funtikova AN, Fito M, Schroder H. Prenatal nutrition and the risk of adult obesity: Long-term effects of nutrition on epigenetic mechanisms regulating gene expression. *J Nutr Biochem* 2017; 39; 1-14.
- 15 Won SB, Kwon YH. Maternal Consumption of Low-Isoflavone Soy Protein Isolate Confers the Increased Predisposition to Alcoholic Liver Injury in Adult Rat Offspring. *Nutrients* 2018; 10.
- 16 Shirpoor A, Minassian S, Salami S, Khadem-Ansari MH, Yeghiazaryan M. Alpha-lipoic acid decreases DNA damage and oxidative stress induced by alcohol in the developing hippocampus and cerebellum of rat. *Cell Physiol Biochem* 2008; 22; 769-76.
- 17 Sousa Coelho IDDd, Lapa Neto CJC, Souza TGdS, Silva MAd, Chagas CA, Santos KRPd, et al. Protective effect of exogenous melatonin in rats and their offspring on the genotoxic response induced by the chronic consumption of alcohol during pregnancy. *Mutat Res Genet Toxicol Environ Mutagen* 2018; 832-833; 52-60.
- 18 Harnish DC, Malik S, Kilbourne E, Costa R, Karathanasis SK. Control of apolipoprotein AI gene expression through synergistic interactions between hepatocyte nuclear factors 3 and 4. *J Biol Chem* 1996; 271; 13621-8.
- 19 Metzger S, Halaas JL, Breslow JL, Sladek FM. Orphan receptor HNF-4 and bZip protein C/EBP alpha bind to overlapping regions of the apolipoprotein B gene promoter and synergistically activate transcription. *J Biol Chem* 1993; 268; 16831-8.
- 20 Hagan DL, Kienzle B, Jamil H, Hariharan N. Transcriptional regulation of human and hamster microsomal triglyceride transfer protein genes. Cell type-specific

expression and response to metabolic regulators. *J Biol Chem* 1994; 269; 28737-44.

21 Crestani M, Sadeghpour A, Stroup D, Galli G, Chiang JY. Transcriptional activation of the cholesterol 7 $\alpha$ -hydroxylase gene (CYP7A) by nuclear hormone receptors. *J Lipid Res* 1998; 39; 2192-200.

22 Jump DB. Fatty acid regulation of hepatic lipid metabolism. *Curr Opin Clin Nutr Metab Care* 2011; 14; 115-20.

23 Parviz F, Matullo C, Garrison WD, Savatski L, Adamson JW, Ning G, et al. Hepatocyte nuclear factor 4 $\alpha$  controls the development of a hepatic epithelium and liver morphogenesis. *Nat Genet* 2003; 34; 292-6.

24 Duncan SA, Manova K, Chen WS, Hoodless P, Weinstein DC, Bachvarova RF, et al. Expression of transcription factor HNF-4 in the extraembryonic endoderm, gut, and nephrogenic tissue of the developing mouse embryo: HNF-4 is a marker for primary endoderm in the implanting blastocyst. *Proc Natl Acad Sci U S A* 1994; 91; 7598-602.

25 Kyrmizi I, Hatzis P, Katrakili N, Tronche F, Gonzalez FJ, Talianidis I. Plasticity and expanding complexity of the hepatic transcription factor network during liver development. *Genes Dev* 2006; 20; 2293-305.

26 Ladas JA, Hadzopoulou-Cladaras M, Kardassis D, Cardot P, Cheng J, Zannis V, et al. Transcriptional regulation of human apolipoprotein genes ApoB, ApoCIII, and ApoAII by members of the steroid hormone receptor superfamily HNF-4, ARP-1, EAR-2, and EAR-3. *J Biol Chem* 1992; 267; 15849-60.

27 Niehof M, Borlak J. HNF4 $\alpha$  and the Ca-channel TRPC1 are novel disease candidate genes in diabetic nephropathy. *Diabetes* 2008; 57; 1069-77.

28 Higuchi N, Kato M, Tanaka M, Miyazaki M, Takao S, Kohjima M, et al. Effects of insulin resistance and hepatic lipid accumulation on hepatic mRNA expression levels of apoB, MTP and L-FABP in non-alcoholic fatty liver disease. *Exp Ther Med* 2011; 2; 1077-81.

29 Pohl J, Ring A, Korkmaz U, Eehalt R, Stremmel W. FAT/CD36-mediated long-chain fatty acid uptake in adipocytes requires plasma membrane rafts. *Mol Biol Cell* 2005; 16; 24-31.

30 Noshmehr H, D'Amico E, Farilla L, Hui H, Wawrowsky KA, Mlynarski W, et al. Fatty acid translocase

(FAT/CD36) is localized on insulin-containing granules in human pancreatic beta-cells and mediates fatty acid effects on insulin secretion. *Diabetes* 2005; 54; 472-81.

31 Hoosdally SJ, Andress EJ, Wooding C, Martin CA, Linton KJ. The Human Scavenger Receptor CD36: glycosylation status and its role in trafficking and function. *J Biol Chem* 2009; 284; 16277-88.

32 Campbell SE, Tandon NN, Woldegiorgis G, Luiken JJ, Glatz JF, Bonen A. A novel function for fatty acid translocase (FAT)/CD36: involvement in long chain fatty acid transfer into the mitochondria. *J Biol Chem* 2004; 279; 36235-41.

33 Li W, Febbraio M, Reddy SP, Yu DY, Yamamoto M, Silverstein RL. CD36 participates in a signaling pathway that regulates ROS formation in murine VSMCs. *J Clin Invest* 2010; 120; 3996-4006.

34 Krammer J, Digel M, Eehalt F, Stremmel W, Fullekrug J, Eehalt R. Overexpression of CD36 and acyl-CoA synthetases FATP2, FATP4 and ACSL1 increases fatty acid uptake in human hepatoma cells. *Int J Med Sci* 2011; 8; 599-614.

35 Koonen DP, Jacobs RL, Febbraio M, Young ME, Soltys CL, Ong H, et al. Increased hepatic CD36 expression contributes to dyslipidemia associated with diet-induced obesity. *Diabetes* 2007; 56; 2863-71.

36 Rondinone CM, Trevillyan JM, Clampit J, Gum RJ, Berg C, Kroeger P, et al. Protein tyrosine phosphatase 1B reduction regulates adiposity and expression of genes involved in lipogenesis. *Diabetes* 2002; 51; 2405-11.

37 Klaman LD, Boss O, Peroni OD, Kim JK, Martino JL, Zabolotny JM, et al. Increased energy expenditure, decreased adiposity, and tissue-specific insulin sensitivity in protein-tyrosine phosphatase 1B-deficient mice. *Mol Cell Biol* 2000; 20; 5479-89.

38 Elchebly M, Payette P, Michaliszyn E, Cromlish W, Collins S, Loy AL, et al. Increased insulin sensitivity and obesity resistance in mice lacking the protein tyrosine phosphatase-1B gene. *Science* 1999; 283; 1544-8.

39 Yeh MM, Brunt EM. Pathology of nonalcoholic fatty liver disease. *Am J Clin Pathol* 2007; 128; 837-47.

40 Kusari J, Kenner KA, Suh KI, Hill DE, Henry RR. Skeletal muscle protein tyrosine phosphatase activity and



tyrosine phosphatase 1B protein content are associated with insulin action and resistance. *J Clin Invest* 1994; 93; 1156-62.

41 Nalpas B, Vassault A, Le Guillou A, Lesgourgues B, Ferry N, Lacour B, et al. Serum activity of mitochondrial aspartate aminotransferase: a sensitive marker of alcoholism with or without alcoholic hepatitis. *Hepatology* 1984; 4; 893-6.

42 Ludwig S, Kaplowitz N. Effect of pyridoxine deficiency on serum and liver transaminases in experimental liver injury in the rat. *Gastroenterology* 1980; 79; 545-9.

43 Zhou SL, Gordon RE, Bradbury M, Stump D, Kiang CL, Berk PD. Ethanol up-regulates fatty acid uptake and

plasma membrane expression and export of mitochondrial aspartate aminotransferase in HepG2 cells. *Hepatology* 1998; 27; 1064-74.

44 Kishi M, Maeyama S, Iwaba A, Ogata S, Koike J, Uchikoshi T. Hepatic veno-occlusive lesions and other histopathological changes of the liver in severe alcoholic hepatitis--a comparative clinicohistopathological study of autopsy cases. *Alcohol Clin Exp Res* 2000; 24; 74S-80S.

45 Yeh MM, Brunt EM. Pathological features of fatty liver disease. *Gastroenterology* 2014; 147; 754-64.