

The assessment effect of antibiotic as prophylaxis against post-operative complications in elective surgery

Ali Enshaei¹, Leila Chodari^{2,3}, Sahar Nourozi⁴, Naser Khalaji^{2*}

¹ Department of surgery, Urmia University of Medical Sciences, Imam Khomeini hospital, Urmia, Iran

² Department of Physiology, Faculty of Medicine, Urmia University of Medical Sciences, Urmia, Iran

³ Neurophysiology Research Center, Urmia University of Medical Sciences, Urmia, Iran

⁴ Faculty of Medicine, Urmia University of Medical Sciences, Urmia, Iran

*Corresponding authors: Naser Khalaji, Address: Department of Physiology, Faculty of Medicine, Urmia University of Medical Sciences, Urmia, Iran, Email: khalaji.Naser@gmail.com

Zip Code: 5714783734, Tel: +984432780803

Abstract

Background & Aims: Nowadays, complications of post-operative surgery are major problems for hospitals and occupying hospital beds that lead to the human and financial costs to society and the individual. The aim of this study is to use of appropriate antibiotic as prophylaxis to reduce infection and length of stay in hospital, prevention of drug resistance and postoperative complications.

Materials & Methods: The present study was carried out on elective surgical patients, including: cholecystitis, inguinal hernia, thyroidectomy and contracture of burn. In this study, the patients were divided into two groups; receiving Cefazolin as prophylactic antibiotic and not receiving Cefazolin.

Results: The results of this study showed that of 618 patients receiving Cefazolin as prophylaxis only 6 patients had postoperative complications and the average length of stay in hospital was 2 days, while in the not receiving Cefazolin group, there were 23 patients of post-operative complications and the average length of stay in hospital was 3.5 days.

Conclusion: According to the findings, antibiotic Cefazolin as prophylaxis reduced post-operative complications and duration of hospitalization.

Keywords: postoperative complications, Cefazolin, prophylaxis, elective surgery

Received 09 Feb 2018, accepted for publication 24 Apr 2018

Introduction

Postoperative complications are one of the major problems in hospitals, which leads to financial costs to communities and the patient (1). Postoperative complications refer to any deviation from the health during the 30 days after surgery (2). These complications include: site infections, bronchial pneumonia, urinary tract infections and kidney failure,

stroke, pulmonary embolism, myocardial infections and cellulitis (2). The most common postoperative complication is surgical site infection (SSI) (3). SSIs are fairly common, occurring in 2 to 5 percent of surgeries involving incisions. Rates of infection differ according to the type of surgery (4). As many as 500,000 SSIs happen in the United States annually (5). Most SSIs are staph infections. SSI is classified as an infection that

begins at the site of a surgical wound fewer than 30 days after the incision is made (6). SSI Leading to increased disease, increased antibiotic use, re-surgery, prolonged admission and care in special sectors, It also increases mortality in patients (7). Causes and risk factors of surgical site infections include surgery that lasts more than 2 hours, elderly, pollution of the operating environment, weak Surgical Techniques, Smoking, weak immune system, diabetes, inappropriate nutritional status of the patient, etc (8, 9).

Antibiotic is type of antimicrobial substance active against bacteria and is the most important type of antibacterial agent for fighting bacterial infections(10). Antibiotic medications are widely used in the treatment and prevention of such infections. They may either kill or inhibit the growth of bacteria(10). Antibiotic therapy as prophylaxis is one of the important factors in surgical procedures (11). Using correctly antibiotics and completing the prescribed dose are very important, as misuse can cause drug resistance.(12) The use of prophylactic antibiotics in the right time is very important as best time to inject prophylactic antibiotic is 30 minutes before the surgery (2). Some studies reported that usage of prophylactic antibiotic have not any positive effects on SSI (13). In other hand, it is reported that prophylactic antibiotic reduced postoperative complications(14, 15).

Cephalosporins are one of the most important prophylactic antibiotics and have the same mode of action as penicillins, they are classified into four groups or generations based on the antimicrobial activity spectrum (16). First-generation cephalosporins are active predominantly against Gram-positive bacteria, and successive generations have increased activity against Gram-negative bacteria (17). Cefazolin is the first generation of cephalosporins that used to treat cellulitis, urinary tract infections, pneumonia, endocarditis, joint infection, and biliary tract infections (18). It is also used to prevent group B streptococcal disease around the time of delivery and before surgery

and is typically given by injection into a muscle or vein (19).

Given that contradictory effect of antibiotics as prophylaxis and the use of appropriate antibiotics and reduction of postoperative complications are very vital, so the aim of this study was to investigate the effect of Cefazolin as prophylactic antibiotic on postoperative complications.

Materials & Methods

This study is an experimental and prospective clinical trial assessment that was conducted in a 6-month period in the surgical wards of Imam Khomeini Hospital in Urmia. Admitted patients for elective surgery including inguinal hernia, cholecystitis, contrast and chest tibial occlusion were selected. This research was carried out in accordance with the Helsinki Protocol for the care and use of patients. 1236 patients (18 to 60 years old) of any type of surgery were randomly divided into two groups:

- 1- **Receiving Cefazolin** (as prophylactic antibiotic) (group 1): (618 patients include: 222 patients with cholecystitis, 280 patients with inguinal hernia, 68 patients with thyroidectomy and 48 patients with contracture of burn)
- 2- **Not receiving Cefazolin** (group 2): (618 patients include: 222 patients with cholecystitis, 280 patients with inguinal hernia, 68 patients with thyroidectomy and 48 patients with contracture of burn)

All patients had same admission conditions and therapeutic trials and they were hospitalized one day before surgery and CBC and U / A tests were performed for them. After serum therapy, patients in the receiving Cefazolin group received 1 gr of Cefazolin ampoules, half an hour before surgery, while patients in the not receiving Cefazolin, only had serum therapy without

any Cefazolin. During the hospitalization, the vital signs and the location of the ulcer were regularly monitored and at the time of discharge, CBC and U / A tests were performed for patients.

All values were analyzed by 1-way analysis of variance (ANOVA), and the Tukey test was used to compare quantitative data. Values less than 0.05 were considered statistically significant in all cases. Results are expressed as means \pm SEM.

Results

Effects of Cefazolin as prophylactic antibiotic on post-operative complications

Figure 1 shows that the use of Cefazolin in the receiving antibiotic group decreased post-operative

complications compared to the not receiving antibiotic group, but this but this increase was not significant. (P= 0.33)

The number of total patients in the each receiving Cefazolin and not receiving antibiotic groups was 618 that 6 patients from first group (3 cases of cholecystectomy, 2 cases of horny inguinal and 1 case of contrast-sensitive diagnosis) referred to hospital with post-operative infection, that

, while 23 patients from the second group (12 cases of cholecystectomy, 7 cases of horny inguinal, 3 case of contrast-sensitive and 1 case of thyroidectomy diagnosis) referred to hospital with post-operative infection. (Fig 2).




Fig 1. Indicates the number of patients with post-operative complications in Not receiving Cefazolin and receiving Cefazolin groups.




Fig 2. Indicates the number of patients with post-operative complications by separating the type of surgery in Not receiving Cefazolin and receiving Cefazolin groups.

Effects of Cefazolin as prophylactic antibiotic on White Blood Cells (WBC) Count in blood:

Results obtained from blood tests depicted in figure 3. According this Fig, WBC significantly ($P=0.007$)

increased in not receiving Cefazolin group compared to admission time. Also, WBC significantly ($P= 0.047$) was more in the not receiving Cefazolin group when compared with receiving Cefazolin group.




Fig 3. Indicates the Count WBC at the time of admission and in Not receiving Cefazolin and receiving Cefazolin groups. Data are shown as mean ± SEM. * indicated significant change compared with Not receiving Cefazolin group. ** indicated significant change compared with Admission time.

Effects of Cefazolin as prophylactic antibiotic on White Blood Cells (WBC) Count in urine:

Figure 4 illustrates that White Blood Cells (WBC) count in urine significantly ($P= 0.012$) increased in not receiving Cefazolin group compared to admission time.

Also, according to Fig 4, White Blood Cells (WBC) count in urine increased in receiving Cefazolin group compared to admission time but not significantly and decreased (not significantly) in receiving Cefazolin group compared to not receiving Cefazolin group.




Fig 4. Indicates the Count WBC in urine at the time of admission and in Not receiving Cefazolin and receiving Cefazolin groups. Data are shown as mean \pm SEM. ** indicated significant change compared with Admission time.

Effect of Cefazolin as prophylactic antibiotic on degree of fever:

One-way ANOVA showed that degree of fever in receiving Cefazolin and not receiving Cefazolin groups significantly ($P< 0.0001$) increased compared to

admission time (Fig 5). Moreover, Cefazolin prescription in receiving Cefazolin group decreased significantly ($P= 0.002$) degree of fever compared to not receiving Cefazolin group.




Fig 5. Indicates the degree of fever at the time of admission and in Not receiving Cefazolin and receiving Cefazolin groups. Data are shown as mean \pm SEM. * indicated significant change compared with Not receiving Cefazolin group. ** indicated significant change compared with Admission time.

Effect of Cefazolin as prophylactic antibiotic on duration of hospitalization:

Figure 6 illustrates the effect of Cefazolin on duration of hospitalization in the study groups. According to this figure, average length of stay in

hospital significantly decreased in receiving Cefazolin group. As, the average length of stay in hospital was 2 days, while in the Not receiving Cefazolin group, the average length of stay in hospital was 3.5 days.




Fig 6. Indicates the duration of hospitalization in Not receiving Cefazolin and receiving Cefazolin groups. Data are shown as mean \pm SEM. * indicated significant change compared with Not receiving Cefazolin group.

Discussion

This study was conducted in Imam Khomeini Hospital in 1397, results of this study showed that Cefazolin treatment as prophylaxis antibiotic before surgery significantly reduces postoperative infection. Increasing white blood cell count is one of the indicators of post-operative infection (20). Our study showed that the mean of WBC in not receiving Cefazolin significantly increased compared to receiving group that this finding confirms the presence of post-operative infection in this group of patients. In this study post-operative infection in patients who did not receive antibiotics were approximately equal in all types of surgeries, so this finding suggests that site infections can occur in any type of surgery. The presence of more white blood cells in the urine of untreated patients with antibiotics also confirms the infection in these patients. According to findings of this study, the average body

temperature was higher in patients without antibiotic pre-treatment, but body temperature changes were not significant, which could indicate that the infection was not severe. Today, there are still contradictions regarding the timing, duration of antibiotic treatment and the choice the type of antibiotic used in surgery (21). In our study, Cefazolin were used half an hour before surgery, which was satisfactorily similar to previous studies. In the last two decades, we have witnessed a considerable increase in both the number and the severity of infections caused by Gram-positive bacteria such *Staphylococcus aureus* (22). In many institutions, Gram-positive bacteria are the cause of more than 50% of all bloodstream infections (23). Today, first-generation cephalosporins are used as a proper choice for preventing post-operative infections which are relatively non-toxic, inexpensive and easily penetrate into soft tissue and bone (16). A clinical human study

has shown that the use of antibiotics 2 hours before skin cutting has the greatest effect on reducing wound infections (22). Over the past decades, more studies have suggested that antibiotic prescribing should continue after surgery (24). However, therapeutic protocols are different and are used in single dose or multiple dosages after surgery (24). Finally, the results of this study showed that the use of Cefazolin half an hour before surgery significantly reduced postoperative infection.

References

- Altamirano F, Oyarce C, Silva P, Toyos M, Wilson C, Lavandero S, et al. Testosterone induces cardiomyocyte hypertrophy through mammalian target of rapamycin complex 1 pathway. *J Endocrinol* 2009;202(2):299–307.
- Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR. Guideline for prevention of surgical site infection, 1999. Hospital Infection Control Practices Advisory Committee. *Infect Control Hosp Epidemiol* 1999;20(4):250–78; quiz 279–80.
- Malone DL, Genuit T, Tracy JK, Gannon C, Napolitano LM. Surgical site infections: reanalysis of risk factors. *J Surg Res* 2002;103(1):89–95.
- Jones RS, Brown C, Opelka F. Surgeon compensation: “Pay for performance,” the American College of Surgeons National Surgical Quality Improvement Program, the Surgical Care Improvement Program, and other considerations. *Surgery* 2005;138(5):829–36.
- Martone WJ, Nichols RL. Recognition, prevention, surveillance, and management of surgical site infections: introduction to the problem and symposium overview. *Clin Infect Dis* 2001;33 Suppl 2:S67-68.
- Ridgeway S, Wilson J, Charlet A, Kafatos G, Pearson A, Coello R. Infection of the surgical site after arthroplasty of the hip. *J Bone Joint Surg Br* 2005;87(6):844–50.
- Lex DJ, Tóth R, Cserép Z, Breuer T, Sági E, Szatmári A, et al. Postoperative differences between colonization and infection after pediatric cardiac surgery—a propensity matched analysis. *J Cardiothorac Surg* 2013;8:166.
- Panah Y. Neonatal intensive care unit nosocomial bacterial infections. *Tehran Univ Med J* 2008;66(5): 349-54.
- Pellowe C. Managing and leading the infection prevention initiative. *J Nurs Manag* 2007; 15(6): 567-73.
- Russell AD. Biocide use and antibiotic resistance: the relevance of laboratory findings to clinical and environmental situations. *Lancet Infect Dis* 2003;3(12):794–803.
- Radji M, Aini F, Fauziyah S. Evaluation of antibiotic prophylaxis administration at the orthopedic surgery clinic of tertiary hospital in Jakarta, Indonesia. *Asian Pac J Trop Dis* 2014;4(3):190.
- Astagneau P, Rioux C, Golliot F, Brucker G, INCISO Network Study Group. Morbidity and mortality associated with surgical site infections: results from the 1997-1999 INCISO surveillance. *J Hosp Infect* 2001;48(4):267–74.
- Small F, Hofmeyr GJ. Antibiotic prophylaxis for cesarean section. *Cochrane Database Syst Rev* 2002;(3):CD000933.
- Chelmow D, Hennesy M, Evantash EG. Prophylactic antibiotics for non-laboring patients with intact membranes undergoing cesarean delivery: an economic analysis. *Am J Obstet Gynecol* 2004;191(5):1661–5.
- Cunningham F, Leveno K, Bloom S, Spong CY, Dashe J. *Williams obstetrics*, 24e. McGraw-hill; 2014.
- Hellbusch LC, Helzer-Julín M, Doran SE, Leibrock LG, Long DJ, Puccioni MJ, et al. Single-dose vs multiple-dose antibiotic prophylaxis in instrumented lumbar fusion—a prospective study. *Surg Neurol* 2008;70(6):622–7; discussion 627.
- Prescott JF. Beta-lactam antibiotics: cephalosporins. *Antimicrobial therapy in veterinary medicine* 2013;153–73.
- AlBuhairan B, Hind D, Hutchinson A. Antibiotic prophylaxis for wound infections in total joint arthroplasty: a systematic review. *J Bone Joint Surg Br* 2008;90(7):915–9.

19. Tita ATN, Rouse DJ, Blackwell S, Saade GR, Spong CY, Andrews WW. Emerging concepts in antibiotic prophylaxis for cesarean delivery: a systematic review. *Obstet Gynecol* 2009;113(3):675–82.
20. Bilgin YM, van de Watering LMG, Eijnsman L, Versteegh MIM, van Oers MHJ, Brand A. Is increased mortality associated with post-operative infections after leukocytes containing red blood cell transfusions in cardiac surgery? An extended analysis. *Transfus Med* 2007;17(4):304–11.
21. Mastronardi L, Tatta C. Intraoperative antibiotic prophylaxis in clean spinal surgery: a retrospective analysis in a consecutive series of 973 cases. *Surg Neurol* 2004;61(2):129–35; discussion 135.
22. Askarian M, Reza Moravveji A, Assadian O. Prescription of prophylactic antibiotics for neurosurgical procedures in teaching hospitals in Iran. *Am J Infect Control* 2007;35(4):260–2.
23. Edmond MB, Wallace SE, McClish DK, Pfaller MA, Jones RN, Wenzel RP. Nosocomial bloodstream infections in United States hospitals: a three-year analysis. *Clin Infect Dis* 1999;29(2):239–44.
24. Kanayama M, Hashimoto T, Shigenobu K, Oha F, Togawa D. Effective prevention of surgical site infection using a Centers for Disease Control and Prevention guideline-based antimicrobial prophylaxis in lumbar spine surgery. *J Neurosurg Spine* 2007;6(4):327–9.